



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.


Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.


CO ₂	=	44.56	:	44	=	1.013	molecules
CaO	=	28.60	:	56	=	0.510	
FeO	=	14.41	:	72	=	0.200	
MgO	=	13.03	:	40	=	0.325	

100.60

yielding the empiric formula—



also—




Dr. Headden's analysis (loc. cit.) gives—


CaCO ₂	=	50.72	:	100	=	0.5072	molecules
MgCO ₂	=	21.98	:	88	=	0.2497	
FeCO ₂	=	27.29	:	116	=	0.2352	
Insoluble	=	0.22					

100.21

yielding the empiric formula—



also—



Hence, Ankerite occurs at Phoenixville in two varieties:—

At the mine with the ratio	Fe : Mg	=	1 : 1
In the tunnel “ “	Fe : Mg	=	2 : 3

OCTOBER 9.

The President, Dr. RUSCHENBERGER, in the chair.

Thirty members present.

Apparent Discriminative Power in the Selection of Food by a Heliozoon.—Prof. LEIDY remarked that he had on several occasions observed actions in the rhizopods apparently indicating a discriminative power in the selection of food. It was certain that they generally swallowed living algæ and animalculæ, and avoided dead ones. He recently had observed a heliozoon eject an article, which appeared to indicate discriminative power. The heliozoon was *Acanthocystis spinifera*. The genus differs from *Actinophrys* in being provided with siliceous rays in addition to the ordinary soft rays. The former emanate from minute disks, forming, as it were, a sort of flexible armor to the body of the *Acanthocystis*. While examining an individual, a rapidly moving, oval, flagellate infusorium, as it was supposed to be, came into contact with several of the soft rays. The infusorium was paralyzed; it assumed a globular shape and became quiescent. It was gradually drawn towards the body of the heliozoon, which projected its armor to meet it, but quickly withdrew it again, and the heliozoon was pushed off beyond the siliceous rays. The same movements

were repeated, and then the infusorian remained outside the siliceous rays. The objects were examined from time to time for several hours. The infusorium was no more drawn towards the body of the heliozoon. After a time it projected a minute bud, which gradually extended into a tortuous tube, proving the supposed infusorium to be a zoospore. It was finally abandoned by the heliozoon, apparently as if it had been determined not to be its proper food.

On Helminthophaga leucobronchialis (Brewster). — SPENCER TROTTER communicated the fact that last winter, while arranging the collection of warblers (*Sylvicolidæ*) in the museum of the Academy, he had discovered among them a specimen of *Helminthophaga leucobronchialis*. This rare species was described by Mr. Wm. Brewster, of Cambridge, Mass., who procured the first specimen at Newtonville, Mass., in the spring of 1870, a description and plate of which he published in the Bulletin of the Nuttall Ornithological Club for April, 1876. A second specimen was taken near Clifton, Delaware Co., Pa., in May, 1877, by Mr. C. D. Wood, an account of which Mr. Trotter sent to the Bulletin for July, 1877. The one now under consideration is, therefore, the third specimen of this rare species which has been discovered.

When he found it, there was no label attached designating its species, sex, or the locality where it was procured; but on the bottom of the stand was written "J. C., 20 Oct. 1862," and three other words that were much blurred, and which he believed to be "not from Bell." Bell is an Ornithologist of New York. The above indicates that John Cassin (for all the birds he examined have the same J. C. written on the stands or labels) had studied the specimen. The date he supposed to be that of its capture; and it is a curious fact that this specimen should have been procured at least eight years before the one from which the first description was taken. He had compared the specimen carefully with Mr. Brewster's description, also with Mr. Wood's specimen, and with these it agrees almost precisely. As the two former were males, from its similarity to them, he believed this one to be a male also. He took pleasure in adding another rare bird to the many which the Academy already possesses.

OCTOBER 16.

The President, Dr. RUSCHENBERGER, in the chair.

Twenty-five members present.

Mineralogical Notes.—Professor GEO. AUG. KÖNIG placed on record the results of an investigation of a peculiar form of Magnetite from Magnet Cove, Ark. The material was obtained from